



## CLIMATE OUTLOOK FOR 2014 SUMMER SEASON IN SERBIA

May 09<sup>th</sup> 2014

Mean summer seasonal temperatures above the **climatological 1981-2010** average are likely to occur. All three individual months have high probability for warmer-than-average conditions. It should be noted that the highest positive anomaly is predicted in July, which is more likely to be extremely warm.

Uncertainties in regional predictions are larger for precipitation than for temperature. In the northern part of the country, which is a part of the Pannonia Plain, summer season precipitation totals are likely to be below-average. In the rest of Serbia the probabilities for below-, near or above- normal conditions are approximately equal. It must be emphasized that it might be possible that some parts, especially mountainous ones, might locally have near- or above- normal summer season totals, due to episodes of the enhanced convection with high intensity rainfall.

Probability for summer season temperature 2 m (%)	Probability for summer season precipitation (%)
<b>Above normal</b> (10, 30, 60)  in the northern part of the country	<b>Below normal</b> (40, 30, 30)  in the northern part of the country
<b>Above normal</b> (20, 30, 50)  in the rest of Serbia	<b>No predictive signal</b> (33, 34, 33)  in the rest of Serbia

The numbers in brackets from left to the right are related to the values of meteorological parameters for: below-, near- and above- normal.