

The seasonal forecast for summer 2017 in Bulgaria

Ilian Gospodinov

National institute of meteorology and hydrology, Bulgaria

ilian.gospodinov@meteo.bg

26 October 2017, Sofia

1. Verification of the seasonal forecast for summer 2017

Tables 1 and 2 show the regular Bulgarian seasonal forecast for the winter season JJA 2017 issued in March (Month-3), April (Month-2), and May (Month-1) 2017 and for the individual months of the summer season issued back up to 3 months prior to the forecast one. The column “Index” gives the assessment of the month or the season based on real data.

Table 1: Scores of the seasonal forecast of mean seasonal temperature for summer 2017.

Temperature	Forecast			Index	Score		
	Month-1	Month-2	Month-3		Month-1	Month-2	Month-3
June	1	1	1	1.89	3	3	3
July	1	1	1	1.29	4	4	4
August	1	1	1	1.69	3	3	3
Summer	1	1	1	1.93	3	3	3

Table 2: Scores of the seasonal forecast of seasonal amount of precipitation for summer 2017.

Precipitation	Forecast			Index	Score		
	Month-1	Month-2	Month-3		Month-1	Month-2	Month-3
June	-1	-1	0	-0.17	3	3	4
July	-1	-1	0	1.06	0	0	2
August	-1	-1	-1	-0.81	4	4	4
Summer	-1	-1	-1	0.05	2	2	2

In average the forecast for temperature scores 3.25 which is very good. In average the seasonal precipitation amount forecast scores 2.5 which is good.

The national seasonal forecast followed roughly the SEECOF and the MedCOF guidelines and the discussions above apply for the assessment of the regional forecast as far as it concerns Bulgaria. The MedCOF and SEECOF forecasts for summer 2017, for the region of Bulgaria, were for near or above normal temperature and no forecast or near normal precipitation.

Since 2012 the Bulgarian seasonal forecast contains an additional sentence that presents an attempt to predict how the upcoming season or month is expected to compare to the same one from the previous year. For this summer it was said that the season should be similar to summer 2016. The summer of 2016 was warm (1.82) with precipitation near or below normal (-0.94). Summer 2017 therefore was similarly warm (1.93) but with more precipitation (0.05) than summer 2016 although the precipitation was around normal in average for the country and therefore the seasonal outlook was not misleading.




Figure 1: Departure of the seasonal mean temperature from normal (1961-1990) (left) and seasonal amount of precipitation in percent of normal (1961-1990) (right) for summer (June-July-August) 2017.

Figure 1, 2, 3, and 4 show maps of the departure from normal (1961-1990) of the seasonal/monthly mean temperature (left) and the seasonal/monthly amount of precipitation in percent of normal (1961-1990) (right) for the summer season as a whole (Fig. 1) and the individual months of June 2017 (Fig.2), July 2017 (Fig.3), and August 2017 (Fig.4). The maps are regular operational products of the Bulgarian weather service and are therefore given with reference to normal based on the period 1961-1990 as with the WMO recommendations.




Figure 2: Departure of the monthly mean temperature from normal (1961-1990) (left) and monthly amount of precipitation in percent of normal (1961-1990) (right) for June 2017.




Figure 3: Departure of the monthly mean temperature from normal (1961-1990) (left) and monthly amount of precipitation in percent of normal (1961-1990) (right) for July 2017.




Figure 4: Departure of the monthly mean temperature from normal (1961-1990) (left) and monthly amount of precipitation in percent of normal (1961-1990) (right) for August 2017.

2. Extreme events

Around 30 June – 1 July the maximum temperatures soared to extreme levels driven by approaching cold front from northwest. They went above 40°C at some places for example - Ruse, 43.6°C.

Fire weather conditions in the southeast were indicating very high risk for wild fires in late August due to long lasting hot and dry weather (Fig. 5).




Figure 5: Fire weather index for 18 August 2017.

3. Explanations

3.1 Regular seasonal forecasting in Bulgaria


The National institute of meteorology and hydrology (NIMH) is the national weather service of Bulgaria. We have been producing regular seasonal forecast for our country since 2005. It is updated once a month at the end of the month as soon as all forecast materials become available. It is based on subjective analysis of the map products from the numerical climate prediction models of the following centers:

- European center for medium range weather forecast, Reading, UK;
- MetOffice, Exeter, UK;
- National center for environmental prediction, USA;
- International research institute, Columbia University, USA;
- Beijing climate center, China;
- Tokyo climate center, Japan;

and the statistical prediction models of the Italian institute of biometeorology Ibimet and the Tokyo climate center. All these materials are available on the websites of the centers.

3.2 Notation rules

The categories “above normal”, “around normal”, and “below normal” by definition have an equal probability of occurrence of 33.3%. The aim of the seasonal forecast is to favor one or two of the three categories based on the analysis of all available forecast materials and assessment of the evolution of large climate structures for the upcoming months. We consider Bulgaria as a region that is relatively small compared to the spatial uncertainties of the modern seasonal forecasting materials. That is why we give a unique forecast valid for the entire country without detailing for different regions except occasionally and only for the first month based on analysis of the medium range weather forecast. The forecast is summarized in tables with the favored categories in color as follows:



We call “season” any three-month period which corresponds to the way the numerical seasonal forecast products are provided by the centers. However since 2011 the seasonal forecast is published only for the calendar season winter, spring, summer, and autumn.

The regular seasonal forecast is available to the public on the website of the institute though only in Bulgarian language.

3.3 Verification rules

In order to quantify the seasonal forecast in terms of categories below, around, and above normal we do the following. Since we give a unique forecast for the expected category for the entire country we need to have a unique assessment of the category of a given month or season. The assessment of the category is based on data from 20 meteorological stations distributed evenly in the country. The data from each of those 20 stations are analyzed. These are records of mean monthly temperature and monthly amount of precipitation from 1950 to present. The percentiles for below, around, and above normal are found for each station based on the latest possible 30-year period 1980-2009. This period is chosen in order to match the base periods of some if not all of the climate centers producing probability map. This reference period is also more suitable to give monthly or seasonal category that would correspond better to the perception of the public. This should be especially true for the thermal category because of the recent overall warming trend. The months and seasons therefore can be attributed a certain category numbered from -2 (below normal) to +2 (above normal). These numbers for all 20 stations and for each individual month or season are then averaged in order to produce a unique category number for the entire country. The forecast itself is also attributed a number that reflects the forecast category. The numbers are -2 (below normal), -1 (below or around normal), 0 (around normal), +1 (above or around normal), and +2 (above normal). In order to assess the skill of our forecast we find the difference between the forecast and the real category. If it is within ± 0.5 we consider that the forecast is excellent (4), within ± 1.0 – very good (3), within ± 1.5 – good (2), and above it is considered to be poor (0). If there is no given preference to any of the three categories we attribute score (1) reasonable, because at least the forecast is not misleading.

References:

Monthly bulletin of the National institute of meteorology and hydrology, Sofia, Bulgaria. Latest issue available online (<http://www.meteo.bg/sites/storm.cfd.meteo.bg.meteo/files/Bulletin.pdf>) and older issues available on demand.

Seasonal forecast for Bulgaria. Latest issue available online (<http://www.meteo.bg/en/node/58>).