Seasonal outlook for DJF 2020/21 conditions over Israel

Yoav Levi and Itzhak Carmona

Israel Meteorological Service

The Seasonal forecast issued by the Israeli Meteorological Service (IMS) is based on ECMWF system 5 seasonal forecast.

1. Prediction of surface air temperature in the surface (2m above ground level) for DJF 2020/21

Figure 1 presents the probabilities for the 3 surface air temperature (2m above ground) terciles over Israel for winter 2020/21 (DJF 2020/21).




Figure 1: The probabilities for the 3 temperature terciles over Israel for winter 2020/21 (DJF 2020/21).

According to figure 1 there is a preferred category above or equal 40%. The highest probability is in the category of above normal, 67%. The probability to be around normal is 27% and the probability to be below normal is only 6%. Hence, DJF averaged surface air temperature in the surface (2m) is expected to be above the normal, in the upper tercile. The mean ensemble of 2020 relies in the percentile above 99% regarding to the distribution of the climatology period 1981-2010.

The predication of surface air temperature for winter in Israel is based on ECMWF seasonal forecast system 5 Ensemble which contains 51 members for the operative prediction. The prediction is relative to the model climatological 1 period 1981/82-2010/11 which is based on Ensemble of 750 members. The prediction is based on the area of 38 grid points, which include the whole area of Israel which can be seen in figure 2.




Figure 2: The area of 38 grid points which are used in order to make surface air temperature seasonal forecast for DJF season in Israel.

2. Prediction of precipitation for DJF 2020/21

Figure 3 presents the probabilities for the 3 precipitation terciles over Israel for winter 2020/21.




Figure 3: The probabilities for the 3 precipitation terciles over Israel for winter 2020/21.

According to figure 3 there is a preferred category above or equal 40% the category is below normal. The highest probability is in the category of below normal, 57%. The probability to be around normal is 22% and the probability to be above normal is only 21%. Hence, DJF accumulated precipitation is expected to be below the normal. The mean ensemble of 2020 relies in the percentile of 11% regarding to the distribution of the climatology period 1981-2010.

The predication of precipitation for winter in Israel is based on ECMWF seasonal forecast system 5 Ensemble which contains 51 members for the operative prediction. The prediction is relative to the model climatological l period 1981/82-2010/11 which is based on Ensemble of 750 members. The prediction is based on the area of 20 grid points where the averaged observed precipitation is above the of 200 mm isohyet (for

all the rainy season not only DJF). Therefore, we include only the region which is north from beer Sheva city (north from latitude 31.3°N). By GIS maps most of the area which is north to beer Sheva have in average more than 200 mm per rainy season.

In summary: the predication of precipitation in Israel include only the area which is north from Beer Sheva city (north from latitude 31.3°N).

The next figure (figure 4) presents the 20 grids point which we are using in order to make prediction for Israel precipitation in DJF season.




Figure 4: The area of 20 grid points which are used in order to make precipitation seasonal forecast for DJF season in Israel.

Please notice the fact that before season DJF 2017/18, (DJF 2016/17 and before) we used to work on ECWMF seasonal forecast system 4, which was approximately two time less in resolution than system 5, therefore we used only 5 grid points in order to predict the winter seasonal forecast precipitation, instead of 20 grid points which are being used today.